arXiv:1211.4452v1 [hep-ph] 19 Nov 2012

MISC-2012-22

Partial mass-degeneracy and spontaneous CP violation in the lepton sector

Takeshi Araki[*] and Hiroyuki Ishida[†]

Maskawa Institute, Kyoto Sangyo University,

Motoyama, Kamigamo, Kita-Ku, Kyoto 603-8555, Japan

Abstract

Inspired by the small squared-mass difference measured in solar neutrino oscillation experiments and observability in neutrinoless double beta decay experiments, we suggest that the partial mass-degenerate limit, in which masses of the first two generation fermions are degenerate, may be a good starting point for understanding the observed fermion mass spectra and mixing patterns. The limit indicates an $O(2)$ symmetry acting on flavor space of the first two generations. We propose a simple model for the lepton sector based on the D_N group, which is a discrete subgroup of $O(2)$, and show that the model can successfully reproduce the experimental data without tuning dimension couplings. We also show that the electron mass and $\theta_{13}^{\rm PMNS}$, which is the smallest mixing angles in the PMNS leptonic mixing matrix, are moderately related with each other through CP violating phases.

---

[*] araki@cc.kyoto-su.ac.jp

[†] h_ishida@cc.kyoto-su.ac.jp

I. INTRODUCTION AND MOTIVATION

Although the Standard Model (SM) of particle physics agrees very well with results from various experiments, some problems and unsatisfactory points have also been pointed out. One of them is the lack of a guiding principle ruling the flavor structure of fermions. In this regard, the introduction of flavor symmetries is one of the most conceivable extensions of the SM to understand the observed fermion mass spectra and flavor mixing patterns and also well motivated from a viewpoint of string theories [1].

Since, as concluded in Ref. [2], most of the flavor symmetries need to be broken either spontaneously or explicitly at energy scales much above the electroweak scale, it is usually no easy task to answer the question, "What kind of symmetries should we introduce?" In order to answer this question, we need to find out the remnants of such broken symmetries in a low energy theory, and small parameters could be important for this purpose. If the symmetry breaking was slight, their breaking terms should correspondingly be small. Moreover, even if they were grossly broken, the breaking effects might be realized in low energies as effective non-renormalizable operators suppressed by their breaking scales. For instance, it is well-known that a Majorana neutrino mass term can be constructed with only the SM particle contents at mass-dimension five [3]:

$$\mathcal{L}_{\rm eff} = \frac{f_{ij}}{\Lambda_\nu} L_i L_j H H, \tag{1}$$

where L and H represent the left-handed lepton doublets and the SM Higgs, respectively, and that the lepton number conservation is restored when one sets the neutrino mass term to zero[1]. In this case, the tiny neutrino masses could be regarded as breaking terms of the lepton number symmetry broken at the high energy scale Λ_ν. Hence, it might be said that small parameters in a low energy theory are the manifestation of new symmetries in high energies; zero limits of the small parameters correspond to unbroken limits of the associated symmetries.

In the case of flavor symmetries, their remnants should appear in the fermion mass and flavor mixing terms. For example, one may line up the following candidates:

1. $\theta_{13}^{\rm PMNS} \ll \theta_{12}^{\rm PMNS},\ \theta_{23}^{\rm PMNS}$ (or $V_{e3}^{\rm PMNS} \ll$ the others),

---

[1] In Ref. [4], the authors relate the tiny neutrino masses with the Peccei-Quinn symmetry.

2. $|\theta_{23}^{\rm PMNS} - 45^\circ| \ll \theta_{23}^{\rm PMNS}$ (or $|V_{\mu 3}^{\rm PMNS} - 1/\sqrt{2}| \ll V_{\mu 3}^{\rm PMNS}$),
3. $\Delta m_{12}^2 = (m_2^\nu)^2 - (m_1^\nu)^2 \ll \Delta m_{23}^2 = |(m_3^\nu)^2 - (m_2^\nu)^2|$,
4. $m_1^{u,d,\ell},\ m_2^{u,d,\ell} \ll m_3^{u,d,\ell}$,
5. $\theta_{ij}^{\rm CKM} \ll \theta_{ij}^{\rm PMNS}$ (or $V_{ij}^{\rm CKM} \ll V_{ij}^{\rm PMNS}$),

where $\theta_{ij}^{\rm PMNS}$ and $\theta_{ij}^{\rm CKM}$ stand for the mixing angles included in the Pontecorvo-Maki-Nakagawa-Sakata (PMNS), $V^{\rm PMNS}$, and Cabibbo-Kobayashi-Maskawa (CKM), $V^{\rm CKM}$, mixing matrices[2], respectively, and m_i^f with $f = u, d, \ell, \nu$ and $i = 1, 2, 3$ denote masses of the up-type quarks, down-type quarks, charged leptons and neutrinos. The first two items have attracted a lot of attention over the years as they indicate the limit of $\theta_{13}^{\rm PMNS} = 0^\circ$ and $\theta_{23}^{\rm PMNS} = 45^\circ$, suggesting the μ-τ permutation symmetry [6] or non-Abelian discrete flavor symmetries [7] in the neutrino sector. In fact a number of models based on these symmetries have been proposed [8]. The recent reactor [9] and long-baseline [10] neutrino oscillation experiments, however, disfavor the vanishing $\theta_{13}^{\rm PMNS}$: the latest data of the DAYA-BAY experiment [11] give us 7.7σ deviations from the vanishing $\theta_{13}^{\rm PMNS}$ with a rather large central value of $\theta_{13}^{\rm PMNS} \simeq 8.7^\circ$. Furthermore, considerable deviations of $\theta_{23}^{\rm PMNS}$ from 45° have also been found in the global analysis of neutrino oscillation experiments [12–14], yet its confidence level is not high enough to conclude.

In this work, instead of these two, we would turn our attention to the remaining three items listed above and discuss their consequences. The third item turns out quasi-mass-degeneracy between the first and second generation neutrinos unless m_1^ν is much smaller than m_2^ν in the normal neutrino mass ordering case. Let us focus on this partial quasi-degenerate region (roughly $m_2^\nu = 0.05 \sim 0.10$ eV) and consider the effective Majorana neutrino mass operator in Eq. (1). In the limit of $m_1^\nu = m_2^\nu$, the Majorana neutrino mass matrix comes to respect the $O(2)$ flavor symmetry [15]

$$R^T \begin{pmatrix} m_1^\nu & 0 & 0 \\ 0 & m_1^\nu & 0 \\ 0 & 0 & m_3^\nu \end{pmatrix} R = P^T \begin{pmatrix} m_1^\nu & 0 & 0 \\ 0 & m_1^\nu & 0 \\ 0 & 0 & m_3^\nu \end{pmatrix} P = \begin{pmatrix} m_1^\nu & 0 & 0 \\ 0 & m_1^\nu & 0 \\ 0 & 0 & m_3^\nu \end{pmatrix}, \tag{2}$$

---

[2] In this paper, we adopt the standard parametrization defined by the Particle Data Group [5].

where

$$R = \begin{pmatrix} \cos\theta & \sin\theta & 0 \\ -\sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad P = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \tag{3}$$

In other words, L_1 and L_2 belong to a doublet representation of $O(2)$, e.g. $\mathbf{2}_n$, while L_3 behaves as a singlet representation, e.g. $\mathbf{1}$. (Notations of $O(2)$ are defined in Appendix.) The observed slight mass splitting between m_1^ν and m_2^ν is regarded as a breaking term of the $O(2)$ symmetry. At first glance, this partial degenerate limit seems to conflict with the charged lepton masses because they are strongly hierarchical. In the case of Dirac mass terms, however, one can assign a different doublet representation, $\mathbf{2}_{m\neq n}$, or the singlet representation $\mathbf{1}$ to the right-handed charged leptons, leading to

$$M^\ell = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & m_3^\ell \end{pmatrix} \quad \text{or} \quad M^\ell = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ m_{31} & m_{32} & m_{33} \end{pmatrix}, \tag{4}$$

respectively, and one can see that the electron and muon are degenerate with zero mass in both cases. In this sense, the partial degenerate limit is applicable to not only the charged lepton sector but also the quark sectors. Rather, this picture appears reasonable since we can relate smallness of the light charged fermion masses with the $O(2)$ symmetry breaking. Also, we may be able to naturally understand some phenomenological relations among the CKM matrix elements [16].

The $O(2)$ symmetry breaking triggers flavor mixing at the same time. On one hand, in the quark and charged lepton sectors, small flavor mixings are expected because the breaking terms are supposed to be much smaller than the leading terms defined in Eq. (4), and thus the observed small CKM mixing could be obtained[3]. On the other hand, in the neutrino sector, flavor mixing can be large since the leading neutrino mass matrix in Eq. (2) is almost proportional to the unit matrix if the neutrino mass hierarchy is mild. Then, it may be possible to realize the large PMNS mixing even thought the breaking terms are small.

---

[3] We notice that mixing between the first and second generations is not necessarily small because of the mass degeneracy. This could explain reason why $\theta_{12}^{\rm CKM}$ is a little larger than the others



FIG. 1: The effective mass, $\langle m_{ee} \rangle$, of neutrinoless double beta decay as a function of m_2^ν within 3σ errors of the oscillation parameters from [13] while varying CP violating phases from 0 to 2π. The current upper bound by combining results from the EXO [17] and KamLAND-Zen [18] experiments at 90% C.L. and the expected future bound by the next generation EXO and KamLAND-Zen experiments [19] are also shown.

To summarize, the partial degenerate limit seems to fit the observed fermion mass spectra and mixing patterns, and moreover it suggests an $O(2)$ flavor symmetry (or its subgroups) acting on flavor space of the first two generations. Yet another motivation to look at the partial quasi-degenerate region of neutrino masses is that the effective mass, $\langle m_{ee} \rangle$, of neutrinoless double beta decay is not vanishing even in the case of normal neutrino mass ordering[4]. In Fig. 1, we plot $\langle m_{ee} \rangle$ as a function of m_2^ν within 3σ errors of the neutrino oscillation parameters from [13] while varying CP phases from 0 to 2π. For example, if $m_2^\nu > 0.05$ eV, then $\langle m_{ee} \rangle > 0.01$ eV, which would be accessible by the next generation EXO and KamLAND-Zen experiments [19]. Therefore, the partial quasi-degenerate region can be confirmed or excluded in the near future if the neutrinos are Majorana particles.

In the next section, we show a simple model for the lepton sector by means of the D_N

---

[4] We would like to thank E. Takasugi for making us aware of this point.

symmetry, which is one of discrete subgroups of $O(2)$, where the order of D_N is supposed to be sufficiently large so that all the doublet representations used in our model obey the same tensor product rules as those of $O(2)$. In this case, we are free from massless Nambu-Goldstone bosons, gauge anomalies etc. and thus can concentrate on the flavor puzzles. Inclusion of the quark sectors and enlargement of the symmetry to $O(2)$ will be studies elsewhere. Section III is devoted to summary and discussions. Notations and tensor product rules of $O(2)$ are summarized in Appendix.

II. MODEL

A. masses and mixings

	L_I	L_3	ℓ_i	H	S_I
D_N	$\mathbf{2}_2$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{1}$	$\mathbf{2}_1$

TABLE I: Particle contents and charge assignments of the model, where $I = 1, 2$ and $i = 1 \cdots 3$ denote the indices of generations, L and ℓ represent the left- and right-handed leptons, and H and $S_{1,2}$ are the SM Higgs and gauge singlet scalar fields, respectively.

We embed the first two generations of the left-handed leptons into the doublet representation $\mathbf{2}_2$, while we assign the trivial singlet representation $\mathbf{1}$ to the other leptons and the SM Higgs[5], giving rise to the partial mass-degenerate limit. In order to lift the mass degeneracy, the D_N flavor symmetry must be broken by a doublet representation, so that we introduce a set of gauge singlet scalars $S_{1,2}$ belonging to $\mathbf{2}_1$. Particle contents and charge assignments of the model are summaries in Table I. We consider the effective Majorana neutrino mass operator in Eq. (1) so as to keep our discussions as general as possible. Under the D_N flavor symmetry, the charged lepton Yukawa and Majorana neutrino mass terms are given by

---

[5] Alternatively, one can embed ℓ_1 and ℓ_2 into a doublet representation, too, as mentioned in Introduction. In this case, however, the electron and muon masses tend to be close to each other, so that we do not consider this assignment in this work.

$$\begin{aligned}\mathcal{L}_f \;=\; & y_i^0\, \overline{L}_3 H \ell_i + \frac{y_i}{\Lambda_F^2}\overline{L}_I H \ell_i (S^2)_I + \frac{y_i^{'}}{\Lambda_F^2}\overline{L}_I H \ell_i (S^{*2})_I + \frac{y_i^{''}}{\Lambda_F^2}\overline{L}_I H \ell_i (|S|^2)_I \\ & + \frac{f_\nu}{\Lambda_\nu} L_I L_I H H + \frac{f_\nu'}{\Lambda_\nu} L_3 L_3 H H \\ & + \frac{g_\nu}{\Lambda_\nu \Lambda_F^2} L_3 L_I H H (S^2)_I + \frac{g_\nu'}{\Lambda_\nu \Lambda_F^2} L_3 L_I H H (S^{*2})_I + \frac{g_\nu^{''}}{\Lambda_\nu \Lambda_F^2} L_3 L_I H H (|S|^2)_I \\ & + \frac{h_\nu}{\Lambda_\nu \Lambda_F^4} (L_J L_K)_I H H (S^4)_I + \frac{h_\nu'}{\Lambda_\nu \Lambda_F^4} (L_J L_K)_I H H (S^{*4})_I + \frac{h_\nu^{''}}{\Lambda_\nu \Lambda_F^4} (L_J L_K)_I H H (|S|^4)_I \\ & + \frac{h_\nu^{'''}}{\Lambda_\nu \Lambda_F^4} (L_J L_K)_I H H (S^2 |S|^2)_I + \frac{h_\nu^{''''}}{\Lambda_\nu \Lambda_F^4} (L_J L_K)_I H H (S^{*2} |S|^2)_I, \end{aligned} \tag{5}$$

where y_i, f_ν, g_ν and h_ν are dimensionless couplings, Λ_F describes a breaking scale of the D_N flavor symmetry and we have ignored the next-to-next-to-leading terms in the charged lepton sector. Note that the gauge singlet scalars S_I are postulated to be complex fields, and we invoke spontaneous CP violation (SCPV) via complex vacuum expectation values (VEVs) of S_I; otherwise the electron remains massless within the given particle contents even if one takes into account further higher order terms. We shall discuss the availability of SCPV in the next subsection and here define VEVs of the scalars as

$$\langle H \rangle = v, \quad \langle S_I \rangle = (s_1 e^{i\phi_1} \quad s_2 e^{i\phi_2})^T. \tag{6}$$

After the scalars acquire the VEVs, the charged lepton mass matrix takes the form of

$$\begin{aligned}\frac{1}{v} M^\ell \;=\; & \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ y_1^0 & y_2^0 & y_3^0 \end{pmatrix} \\ & + \frac{1}{\Lambda_F^2}\left[\begin{pmatrix} y_1\delta_1 & y_2\delta_1 & y_3\delta_1 \\ y_1\delta_2 & y_2\delta_2 & y_3\delta_2 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} y_1'\delta_1^* & y_2'\delta_1^* & y_3'\delta_1^* \\ y_1'\delta_2^* & y_2'\delta_2^* & y_3'\delta_2^* \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} y_1^{''}\delta_a & y_2^{''}\delta_a & y_3^{''}\delta_a \\ y_1^{''}\delta_b & y_2^{''}\delta_b & y_3^{''}\delta_b \\ 0 & 0 & 0 \end{pmatrix} \right], \end{aligned} \tag{7}$$

where

$$\delta_1 = (s_1^2 e^{2i\phi_1} - s_2^2 e^{2i\phi_2}), \quad \delta_2 = 2 s_1 s_2 e^{i(\phi_1 + \phi_2)}, \tag{8}$$

$$\delta_a = s_1^2 - s_2^2, \quad \delta_b = 2 s_1 s_2 \cos(\phi_1 - \phi_2), \tag{9}$$



FIG. 2: $\sin^2\theta_{13}^{\rm PMNS}$ as a function of $\sin^2\theta_{23}^{\rm PMNS}$ for the cases of normal (left-panel) and inverted (right-panel) neutrino mass orderings. The 1σ error bars are also plotted for the normal and inverted cases from Ref. [13] and Ref. [12], respectively.

while the neutrino mass matrix is given as

$$\frac{\Lambda_\nu}{v^2}\mathcal{M}_\nu = \begin{pmatrix} f_\nu & 0 & 0 \\ 0 & f_\nu & 0 \\ 0 & 0 & f'_\nu \end{pmatrix} + \frac{1}{\Lambda_F^2}\begin{pmatrix} 0 & 0 & g_\nu\delta_1 + g'_\nu\delta_1^* + g''_\nu\delta_a \\ 0 & 0 & g_\nu\delta_2 + g'_\nu\delta_2^* + g''_\nu\delta_b \\ g_\nu\delta_1 + g'_\nu\delta_1^* + g''_\nu\delta_a & g_\nu\delta_2 + g'_\nu\delta_2^* + g''_\nu\delta_b & 0 \end{pmatrix}$$

$$+\frac{1}{\Lambda_F^4}\begin{pmatrix} h_\nu\epsilon_1 + h'_\nu\epsilon_1^* + h''_\nu\epsilon_a + h'''_\nu\epsilon_x + h''''_\nu\epsilon_x^* & h_\nu\epsilon_2 + h'_\nu\epsilon_2^* + h''_\nu\epsilon_b + h'''_\nu\epsilon_y + h''''_\nu\epsilon_y^* & 0 \\ h_\nu\epsilon_2 + h'_\nu\epsilon_2^* + h''_\nu\epsilon_b + h'''_\nu\epsilon_y + h''''_\nu\epsilon_y^* & -(h_\nu\epsilon_1 + h'_\nu\epsilon_1^* + h''_\nu\epsilon_a + h'''_\nu\epsilon_x + h''''_\nu\epsilon_x^*) & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (10)$$

with

$$\epsilon_1 = \delta_1^2 - \delta_2^2, \quad \epsilon_2 = 2\delta_1\delta_2, \quad (11)$$

$$\epsilon_a = \delta_a^2 - \delta_b^2, \quad \epsilon_b = 2\delta_a\delta_b, \quad (12)$$

$$\epsilon_x = \delta_1\delta_a - \delta_2\delta_b, \quad \epsilon_y = \delta_1\delta_b + \delta_2\delta_a. \quad (13)$$

Obviously, there exist a sufficient number of parameters to fit the experimental data, and there are no predictions for the fermion masses and mixings. Nevertheless, we would emphasize that the model can reproduce experimental data without making the dimensionless couplings hierarchical, thanks to the Froggatt-Nielsen [20] like mechanism. For instance, in Fig 2, we choose the following parameter spaces:

$$\begin{aligned}
&y_3^0 = 1.0,\ y_1^0 = y_2^0 = 1.2,\ y_1 = -y_2 = y_3 = y_1' = y_2' = -y_3' = 0.8, \\
&y_1^{''} = -y_2^{''} = -y_3^{''} = 0.8 \sim 1.3, \qquad (14) \\
&f_\nu' = 1.0,\ f_\nu = 0.90 \sim 0.95,\ g_\nu = g_\nu^{''} = 0.9,\ g_\nu' = 0.8 \sim 1.3, \\
&h_\nu = h_\nu' = h_\nu^{''} = -h_\nu^{'''} = -h_\nu^{''''} = 0.8 \sim 1.3, \qquad (15) \\
&\frac{s_{1,2}}{\Lambda_F} = 0.15 \sim 0.30,\ \phi_{1,2} = 0 \sim 2\pi, \qquad (16)
\end{aligned}$$

for the case of normal neutrino mass ordering and

$$\begin{aligned}
&y_3^0 = 1.0,\ y_1^0 = y_2^0 = 1.2,\ y_1 = -y_2 = y_3 = y_1' = y_2' = -y_3' = 0.8, \\
&y_1^{''} = -y_2^{''} = -y_3^{''} = 0.8 \sim 1.3, \qquad (17) \\
&f_\nu' = 1.0,\ f_\nu = 1.05 \sim 1.10,\ g_\nu = g_\nu^{''} = -0.9,\ g_\nu' = 1.0 \sim 1.5, \\
&h_\nu = h_\nu' = h_\nu^{''} = -h_\nu^{'''} = -h_\nu^{''''} = -(1.0 \sim 1.5), \qquad (18) \\
&\frac{s_{1,2}}{\Lambda_F} = 0.15 \sim 0.30,\ \phi_{1,2} = 0 \sim 2\pi, \qquad (19)
\end{aligned}$$

for the inverted case, and numerically compute $\sin^2\theta_{23}^{\rm PMNS}$ and $\sin^2\theta_{13}^{\rm PMNS}$ while imposing the 1σ constraints

$$\frac{\Delta m_{12}^2}{\Delta m_{23}^2} = \begin{cases} (2.94 \sim 3.35)\times 10^{-2} & \text{for Normal} \\ (2.97 \sim 3.30)\times 10^{-2} & \text{for Inverted} \end{cases}, \qquad (20)$$

$$\sin^2\theta_{12}^{\rm PMNS} = \begin{cases} 0.291 \sim 0.325 & \text{for Normal} \\ 0.303 \sim 0.336 & \text{for Inverted} \end{cases}, \qquad (21)$$

from Refs. [13] (for Normal) and [12] (for Inverted) and the charged lepton mass ratios at the Z-boson mass scale:

$$\frac{m_e}{m_\mu} = 4.74\times 10^{-3},\ \frac{m_\mu}{m_\tau} = 5.88\times 10^{-2}, \qquad (22)$$

from Ref. [21]. As can be seen from the figures, a nearly maximal $\theta_{23}^{\rm PMNS}$ and relatively large $\theta_{13}^{\rm PMNS}$ are successfully obtained. In Fig. 3, we further place 1σ constraints on $\sin^2\theta_{23}^{\rm PMNS}$ and $\sin^2\theta_{13}^{\rm PMNS}$ and plot s_1/Λ_F versus s_2/Λ_F. Although this analysis is based on the specific parameter choices, the figures show that $s_{1,2}/\Lambda_F$ typically run from 0.18 to 0.28 only if strong hierarchies for the dimensionless couplings are not assumed. Furthermore,



FIG. 3: s_2/Λ_F versus s_1/Λ_F for the cases of normal (left-panel) and inverted (right-panel) neutrino mass orderings. All the 1σ constraints of neutrino oscillation parameters are placed.

in these parameter spaces, the second generation neutrino mass and the effective mass of neutrinoless double beta decay are estimated as

$$m_2^\nu = 0.07 \sim 0.08 \text{ eV}, \quad \langle m_{ee} \rangle = 0.07 \sim 0.08 \text{ eV}, \tag{23}$$

for both normal and inverted mass orderings.

Let us comment on a moderate correlation between the electron mass and $\theta_{13}^{\rm PMNS}$ through the CP violating phases $\phi_{1,2}$. In the model under consideration, the limit of $\phi_1 = \phi_2$ corresponds to the massless limit of the electron. This means that there exists a novel relation between the electron mass and CP violation like the Timeon model [22] and that ϕ_1 needs to be nearly equal to ϕ_2. Suppose $\phi_2 = \phi_1 + \delta\phi$ and $\delta\phi \ll 1$, then the first and second rows of the charged lepton mass matrix can be rewritten as

$$\begin{pmatrix} \left[y_1 e^{2i\phi_1} + y_1' e^{-2i\phi_1} + y_1''\right](s_1^2 - s_2^2) & \cdots & \cdots \\ \left[y_1 e^{2i\phi_1} + y_1' e^{-2i\phi_1} + y_1''\right] 2 s_1 s_2 & \cdots & \cdots \\ 0 & \cdots & \cdots \end{pmatrix}$$
$$+ i\delta\phi \begin{pmatrix} -\left[y_1 e^{2i\phi_1} - y_1' e^{-2i\phi_1}\right] 2 s_2^2 & \cdots & \cdots \\ \left[y_1 e^{2i\phi_1} - y_1' e^{-2i\phi_1}\right] 2 s_1 s_2 & \cdots & \cdots \\ 0 & \cdots & \cdots \end{pmatrix}, \tag{24}$$

where each term possesses rank $= 1$. The first term is mainly responsible for the muon mass while the second one is for the electron mass as long as y_i and ϕ_1 are not tuned.

Therefore, in the first term, the first row should be somewhat smaller than the second row, which requires $s_1 \simeq s_2$. Meanwhile, a 1-3 element of the neutrino mass matrix can similarly be rewritten as

$$(\mathcal{M}_\nu)_{13} \propto \left[g_\nu e^{2i\phi_1} + g'_\nu e^{-2i\phi_1} + g''_\nu\right](s_1^2 - s_2^2) - i\delta\phi\left[g_\nu e^{2i\phi_1} - g'_\nu e^{-2i\phi_1}\right] 2s_2^2. \tag{25}$$

Since $(\mathcal{M}_\nu)_{13}$ is a dominant factor in determining the size of $\theta_{13}^{\rm PMNS}$, $s_1^2 - s_2^2 \ll 1$ and $\delta\phi \ll 1$ result in a small $\theta_{13}^{\rm PMNS}$.

B. scalar potential and spontaneous CP violation

In this section, we discuss the possibility for SCPV in our model. We expand the singlet scalar fields with its VEVs as

$$S = \begin{pmatrix} S_1 \\ S_2 \end{pmatrix} \to \begin{pmatrix} s_1 e^{i\phi_1} + S_1 \\ s_2 e^{i\phi_2} + S_2 \end{pmatrix}. \tag{26}$$

We write the full scalar potential upto renormalizable level:

$$V = V_H + V_S + V_{HS}, \tag{27}$$

$$V_H = \alpha\left|H\right|^2 + \beta\left|H\right|^4, \tag{28}$$

$$\begin{aligned} V_S = {} & \alpha_S\left|S\right|^2 + \alpha'_S{\rm Re}\left[S^2\right] + \beta_S^a\left|S\right|^2_{\mathbf{1}}\left|S\right|^2_{\mathbf{1}} + \beta_S^b\left|S\right|^2_{\mathbf{1}'}\left|S\right|^2_{\mathbf{1}'} + \beta_S^c\left|S\right|^2_{\mathbf{2}}\left|S\right|^2_{\mathbf{2}} \\ & + \beta'_S{\rm Re}\left[S^4\right] + \gamma_S\left|S\right|^2{\rm Re}\left[S^2\right], \end{aligned} \tag{29}$$

$$V_{HS} = \lambda\left|H\right|^2\left|S\right|^2 + \lambda'\left|H\right|^2{\rm Re}\left[S^2\right], \tag{30}$$

where the couplings $\beta_S^A\,(A = a, b, c)$ distinguish different combinations of $S_{1,2}$ under the D_N tensor product rules, and all of the couplings are supposed to be real. Substituting the VEVs into the potential, we obtain

$$V_H = \alpha v^2 + \beta v^4, \tag{31}$$

$$\begin{aligned} V_S = {} & \alpha_S(s_1^2 + s_2^2) + \alpha'_S(s_1^2\cos 2\phi_1 + s_2^2\cos 2\phi_2) + \beta_S^a(s_1^2 + s_2^2)^2 \\ & -4\beta_S^b s_1^2 s_2^2 \sin^2(\phi_1 - \phi_2) + \beta_S^c\{(s_1^2 - s_2^2)^2 + 4s_1^2 s_2^2\cos^2(\phi_1 - \phi_2)\} \\ & +\beta'_S\left[s_1^4\cos 4\phi_1 + s_2^4\cos 4\phi_2 + 2s_1^2 s_2^2\cos[2(\phi_1 + \phi_2)]\right] \\ & +\gamma_S(s_1^2 + s_2^2)(s_1^2\cos 2\phi_1 + s_2^2\cos 2\phi_2), \end{aligned} \tag{32}$$

$$V_{HS} = \lambda v^2(s_1^2 + s_2^2) + \lambda' v^2(s_1^2\cos 2\phi_1 + s_2^2\cos 2\phi_2). \tag{33}$$

Hereafter, we presume that the singlet scalar fields were completely decoupled from the theory at a high energy scale and investigate only the potential V_S. Then, the minimization conditions are calculated as

$$\begin{aligned}
\frac{\partial V_S}{\partial \phi_1} &= -2s_1^2 \left[\ \left\{\alpha_S' + \gamma_S \left(s_1^2 + s_2^2\right)\right\} \sin 2\phi_1 + 2(\beta_S^b + \beta_S^c) s_2^2 \sin\left[2\left(\phi_1 - \phi_2\right)\right] \right. \\
&\qquad \left. + 2\beta_S' \left(s_1^2 \sin 4\phi_1 + s_2^2 \sin\left[2\left(\phi_1 + \phi_2\right)\right]\right)\ \right] = 0\,, \qquad (34) \\
\frac{\partial V_S}{\partial \phi_2} &= -2s_2^2 \left[\ \left\{\alpha_S' + \gamma_S \left(s_1^2 + s_2^2\right)\right\} \sin 2\phi_2 - 2(\beta_S^b + \beta_S^c) s_1^2 \sin\left[2\left(\phi_1 - \phi_2\right)\right]\right] \\
&\qquad + 2\beta_S' \left(s_2^2 \sin 4\phi_2 + s_1^2 \sin\left[2\left(\phi_1 + \phi_2\right)\right]\right)\ \big] = 0\,, \qquad (35) \\
\frac{\partial V_S}{\partial s_1} &= 2s_1 \left[\ \alpha_S + \alpha_S' \cos 2\phi_1 + 2\beta_S^a (s_1^2 + s_2^2) - 4\beta_S^b s_2^2 \sin^2(\phi_1 - \phi_2) \right. \\
&\qquad + 2\beta_S^c \{s_1^2 - s_2^2 + 2s_2^2 \cos^2(\phi_1 - \phi_2)\} + 2\beta_S' (s_1^2 \cos 4\phi_1 + s_2^2 \cos\left[2(\phi_1 + \phi_2)\right]) \\
&\qquad \left. + \gamma_S \{s_2^2 \cos 2\phi_2 + (2s_1^2 + s_2^2) \cos 2\phi_1\}\ \right] = 0\,, \qquad (36) \\
\frac{\partial V_S}{\partial s_2} &= 2s_2 \left[\ \alpha_S + \alpha_S' \cos 2\phi_2 + 2\beta_S^a (s_1^2 + s_2^2) - 4\beta_S^b s_1^2 \sin^2(\phi_1 - \phi_2) \right. \\
&\qquad - 2\beta_S^c \{s_1^2 - s_2^2 - 2s_1^2 \cos^2(\phi_1 - \phi_2)\} + 2\beta_S' (s_2^2 \cos 4\phi_2 + s_1^2 \cos\left[2(\phi_1 + \phi_2)\right]) \\
&\qquad \left. + \gamma_S \{s_1^2 \cos 2\phi_1 + (s_1^2 + 2s_2^2) \cos 2\phi_2\}\ \right] = 0\,. \qquad (37)
\end{aligned}$$

Let us set α_S', β_S' and γ_S to zero just for simplicity, then the first two conditions become

$$2(\beta_S^b + \beta_S^c) s_2^2 \sin[2(\phi_1 - \phi_2)] = 0, \quad 2(\beta_S^b + \beta_S^c) s_1^2 \sin[2(\phi_1 - \phi_2)] = 0. \qquad (38)$$

Suppose $\phi_1 \simeq \phi_2$, which is preferred from a model building point of view, the third and fourth conditions give us

$$s_1^2 + s_2^2 = -\frac{\alpha_S}{2(\beta_S^a + \beta_S^c)}\,. \qquad (39)$$

III. SUMMARY AND FUTURE WORKS

Inspired by $\Delta m_{12}^2 \ll \Delta m_{23}^2$, we focus on a neutrino mass range in which the first two generation neutrinos are quasi-degenerate in mass. In the limit of $m_1^\nu = m_2^\nu$, a Majorana neutrino mass matrix respects the $O(2)$ flavor symmetry: the first two generations constitute a doublet representation while the third one acts as a singlet representation. We suggest that this partial mass-degenerate limit may be a good starting point for understanding not only the observed neutrino mass spectrum but also the charged fermion mass spectra and flavor mixings. We propose a simple model for the lepton sector by means of the D_N symmetry, which is one of discrete subgroups of $O(2)$, and find that the model can successfully reproduce the experimental data without tuning dimensionless couplings. In the model, the electron acquires its mass after CP is spontaneously broken. A novel relation between the electron mass and $\theta_{13}^{\rm PMNS}$ through the CP violating phases $\phi_{1,2}$ is also found.

In the present work, we adopt the D_N symmetry as our flavor symmetry in order to concentrate of the flavor puzzles of fermions. It is, of course, very challenging to enlarge the symmetry to $O(2)$ taking into account a new gauge boson and gauge anomalies. Also, the quark sectors should be included, and we need to check whether the CP phases responsible for the electron mass can simultaneously generate the realistic up- and down-quark masses with the Dirac phase in the CKM matrix. Yet another future work is to postulate a specific neutrino mass generation mechanism and implement the Leptogenesis mechanism. These issues will be studied elsewhere.

ACKNOWLEDGMENTS

The authors would thank to J. Kubo and J. Heeck for useful discussions and comments.

Appendix A: Group theory of $O(2)$

$O(2)$ is the group of rotations and reflections in the two-dimensional plane and composed of two singlet, $\mathbf{1}$ and $\mathbf{1}'$, and an countably infinite set of doublet representations. The doublets can be characterized by means of natural numbers $n \in \mathbb{N}$, such as $\mathbf{2}_{n=1,2,3,\ldots}$. The tensor product rules are defined as

$$\mathbf{1}\otimes\mathbf{1}'=\mathbf{1}', \quad \mathbf{1}'\otimes\mathbf{1}'=\mathbf{1}, \tag{A1}$$

$$\begin{array}{ccccccccc} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \otimes & \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} & = & (x_1y_1+x_2y_2) & \oplus & (x_1y_2-x_2y_1) & \oplus & \begin{pmatrix} x_1y_1-x_2y_2 \\ x_1y_2+x_2y_1 \end{pmatrix} \\ \mathbf{2}_n & \otimes & \mathbf{2}_n & = & \mathbf{1} & \oplus & \mathbf{1}' & \oplus & \mathbf{2}_{2n} \end{array}, \tag{A2}$$

$$\begin{array}{ccccccc} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \otimes & \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} & = & \begin{pmatrix} x_1y_1+x_2y_2 \\ x_1y_2-x_2y_1 \end{pmatrix} & \oplus & \begin{pmatrix} x_1y_1-x_2y_2 \\ x_1y_2+x_2y_1 \end{pmatrix} \\ \mathbf{2}_n & \otimes & \mathbf{2}_{m\neq n} & = & \mathbf{2}_{m-n} & \oplus & \mathbf{2}_{m+n} \end{array}. \tag{A3}$$

Note that, in Ref. [23], $O(2)$ is defined in a different basis.

---

[1] T. Kobayashi, H. P. Nilles, F. Ploger, S. Raby, and M. Ratz, Nucl. Phys. B**768**, 135 (2007); H. P. Nilles, M. Ratz, and P. K.S. Vaudrevange, arXiv:1204.2206 [hep-ph].

[2] C. I. Low and R. R. Volkas, Phys. Rev. D**68**, 033007 (2003); Y. Koide, Phys. Rev. D**71**, 016010 (2005).

[3] S. Weinberg, Phys. Rev. Lett. **43**, 1566 (1979).

[4] C. S. Chen and L. H. Tsai, arXiv:1210.6264 [hep-ph].

[5] J. Beringer *et al.* (Particle Data Group), Phys. Rev. D**86**, 010001 (2012).

[6] T. Fukuyama and H. Nishiura, in *Proceedings of 1997 Shizuoka Workshop on Masses and Mixings of Quarks and Leptons*, hep-ph/9702253; G. Altarelli and F. Feruglio, Phys. Lett. B **439**, 112 (1998); E. Ma and M. Raidal, Phys. Rev. Lett. **87**, 011802 (2001); C. S. Lam, Phys. Lett. B **507**, 214 (2001); K. R. S. Balaji, W. Grimus, and T. Schwets, Phys. Lett. B **508**, 301 (2001).

[7] P. O. Ludl, arXiv:0907.5587 [hep-ph]; H. Ishimori *et al.*, Prog. Theor. Phys. Suppl. **183**, 1 (2010).

[8] G. Altarelli and F. Feruglio, Rev. Mod. Phys. **82**, 2701 (2010); L. Merlo, arXiv:1004.2211 [hep-ph]; C. S. Lam, Phys. Rev. D**83**, 113002 (2011); and references therein.

[9] KamLAND Collaboration, Phys. Rev. D**83**, 052002 (2011); DOUBLE-CHOOZ Collaboration, Phys. Rev. Lett. **108**, 131801 (2012); DAYA-BAY Collaboration, Phys. Rev. Lett. **108**, 171803 (2012); RENO collaboration, Phys. Rev. Lett. **108**, 191802 (2012).

[10] T2K Collaboration, Phys. Rev. Lett. **107**, 041801 (2011); MINOS Collaboration, Phys. Rev. Lett. **107**, 181802 (2011).

[11] DAYA-BAY Collaboration, arXiv:1210.6327 [hep-ex].

[12] D. V. Forero, M. Tórtola, and J. W. F. Valle, Phys. Rev. D**86**, 073012 (2012).

[13] G. L. Fogli *et al*, arXiv:1205.5254 [hep-ph].

[14] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado, and T. Schwetz, arXiv:1209.3023 [hep-ph].

[15] E. I. Lashin, S. Nasri, E. Malkawi, and N. Chamoun, Phys. Rev. D**83**, 013002 (2011); J. Heeck and W, Rodejohann, JHEP **1202**, 094 (2012).

[16] Z. Z. Xing, arXiv:1211.3890.

[17] EXO Collaboration, Phys. Rev. Lett. **109**, 032505 (2012).

[18] KamLAND-Zen Collaboration, Phys. Rev. C**86**, 021601 (2012); arXiv:1211.3863 [hep-ex].

[19] Talks given at *Neutrino 2012, the XXV International Conference on Neutrino Physics and Astrophysics (Kyoto, Japan, 2012)*, website: neu2012.kek.jp.

[20] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. B**147**, 277 (1079).

[21] Z. Z. Xing, H. Zhang, and S. Zhou, Phys. Rev. D**86**, 013013 (2012).

[22] R. Friedberg and T. D. Lee, arXiv:0809.3633 [hep-ph]; T. Araki and C. Q. Geng, Phys. Rev. D**79**, 077301 (2009).

[23] W. Grimus and L. Lavoura, JHEP **0807**, 051 (2008).